

JOURNAL OF ALGEBRA 31, 117–119 (1974)

## Isotopy–Isomorphism Loops of Prime Order\*

ROBERT L. WILSON, JR.<sup>†</sup>*Department of Mathematics, University of Wisconsin, Madison, Wisconsin 53706**Communicated by R. H. Bruck*

Received December 10, 1969

## 1. INTRODUCTION AND DEFINITIONS

(1.1) The main result in this paper is Theorem (2.4) which states that a loop of prime order is isomorphic to all of its loop isotopes if and only if it is a group. The proof is combinatorial and based on a formula due to Bryant and Schneider [2].

(1.2) We say a loop  $(L, \cdot)$  has the *isotopy–isomorphism* property and write  $(L, \cdot) \in I-I$  if  $(L, \cdot)$  is isomorphic to all of its loop isotopes.

We use the following notation from [2].

(1.3) If  $(L, \cdot)$  is a loop and  $a, b \in L$  we denote by  $L(a, b, \circ)$  the principal isotope of  $(L, \cdot)$  whose operation  $\circ$  is defined by

$$x \circ y = (x/b) \cdot (a \setminus y).$$

Any principal loop isotope of  $(L, \cdot)$  (and hence up to isomorphism any loop isotope of  $(L, \cdot)$ ) can be obtained as  $L(a, b, \circ)$  for some choice of  $a, b \in L$ .

(1.4) We write  $L_1 \stackrel{\theta}{\cong} L_2$  if the loop  $L_1$  is isomorphic to  $L_2$  under the mapping  $\theta$ . For a fixed loop  $(L, \cdot)$ , let  $G$  be the set

$$\{\theta \mid L(a, b) \stackrel{\theta}{\cong} L(c, d) \text{ for some } a, b, c, d \in L\}.$$

Bryant and Schneider [2] show that  $G$  with composition forms a group.

\* This note forms part of the author's Ph.D. thesis at the University of Wisconsin. The thesis was written under the direction of Professor Hans Schneider, to whom the author wishes to express great appreciation.

<sup>†</sup> During the research leading to this paper, the author was partially supported under NSF Contract No. 144-7649.

(1.5) Again for a fixed loop  $(L, \cdot)$ , let  $A(a, b)$  denote the group of automorphisms of  $L(a, b, \circ)$ , and  $A = A(1, 1)$ . Note that  $A$  is a subgroup of  $G$  and hence  $|A|$  divides  $|G|$ . For any loop  $(H, *)$ , let  $N_\mu(H, *)$  denote the middle nucleus [1, p. 57] of  $(H, *)$ .

## 2. COUNTING ARGUMENTS

Theorem 5 of [2] shows that the number of principal loop isotopes of a finite loop  $(L, \cdot)$  isomorphic to a particular principal loop isotope  $L(a, b, \circ)$  is

$$\frac{|G| |N_\mu[L(a, b, \circ)]|}{|A(a, b)|}. \quad (2.1)$$

Using the fact [1, p. 57] that the middle nuclei of isotopic loops are isomorphic and hence have the same order, it is immediate that the numerator of (2.1) is actually independent of  $a$  and  $b$ , i.e.,

**PROPOSITION 2.2.** *The number of principal loop isotopes of  $(L, \cdot)$  isomorphic to  $L(a, b, \circ)$  is*

$$\frac{|G| |N_\mu(L, \cdot)|}{|A(a, b)|}.$$

**THEOREM 2.3.** *If  $(L, \cdot)$  is an I-I loop of order  $n$  and  $n^2$  does not divide  $|G|/|A|$ , then  $N_\mu(L, \cdot)$  is not trivial.*

*Proof.* If  $(L, \cdot)$  is an I-I loop, all  $n^2$  of its principal loop isotopes are isomorphic to  $L(1, 1, \cdot) = (L, \cdot)$ . Thus by (2.2),

$$n^2 = \frac{|G|}{|A|} |N_\mu(L, \cdot)|.$$

By hypothesis,  $n^2$  does not divide  $|G|/|A|$  and hence  $|N_\mu(L, \cdot)| > 1$ .

**THEOREM 2.4.** *A loop of prime order is an I-I loop if and only if it is a cyclic group.*

*Proof.* It is known [1, p. 57] that every group is an I-I loop. Thus we need only show that if  $(L, \cdot)$  is an I-I loop of prime order  $p$  then  $(L, \cdot)$  is associative.  $G$  is a subgroup of the symmetric group on  $L$ , and hence  $|G|$  divides  $p!$ . Thus  $|G|/|A|$  divides  $p!$ . Since  $p$  is prime,  $p$  but not  $p^2$  divides  $p!$ . Hence  $p^2$  does not divide  $|G|/|A|$ . By Theorem 2.3,  $N_\mu(L, \cdot)$  is not trivial. For any loop the order of the middle nucleus divides the order of the loop, which in this case is prime, so  $N_\mu(L, \cdot) = L$  and  $L$  is associative.

## 3. NONPRIME ORDERS

In light of Theorem 2.4 one may ask whether the I-I property and associativity are equivalent for any nonprime orders. A construction by the author [3] shows that there is a nonassociative I-I loop for any even order for which there exist nonassociative loops, i.e., even orders  $> 4$ .

## REFERENCES

1. R. H. BRUCK, "A Survey of Binary Systems," Springer-Verlag, Berlin, 1958.
2. B. F. BRYANT AND H. SCHNEIDER, Principal loop-isotopes of quasigroups, *Canad. J. Math.* **18** (1966), 120-125.
3. R. L. WILSON, Quasidirect products of loops, to appear.